

Creation of Locally Compact Topological Space with Calculations

Elena Kotevska and Sonja Chalamani
The Faculty of Technical Sciences at the University st. Kliment
Ohridski 7000 Bitola
Republic of Macedonia
{elena.kotevska@tfb.uklo.edu.mk}
{scalamani@yahoo.com}



ABSTRACT: In this paper, we have demonstrated a way to define groups of cohomologies. We have used illustrations to explain the local compact topological space. We have constructed and also provided correct calculations. We further explained the necessary properties and examined them in detail.

Keywords: Topological Space, Compact Set, p -Measurable Function, p -Measurable Coboundary, Group of Cohomologies

Received: 18 March 2023, Revised 7 May 2023, Accepted 28 June 2023

DOI: 10.6025/jitr/2023/14/3/57-62

Copyright: with authors

1. Introduction

Let X be a topological space, and let G be an Abelian group. In [1] we introduced the Abelian group $\Phi^p(X, G)$ of p -function and we constructed the homeomorphism

$$\delta: \Phi^p(X, G) \rightarrow \Phi^{p+1}(X, G).$$

This homomorphism is defined for every $p \geq 0$ such that for $\varphi \in \Phi^p(X, G)$ the corresponding $\delta\varphi \in \Phi^{p+1}(X, G)$ is given by

$$\begin{aligned} & (\delta\varphi)(x_0, x_1, \dots, x_p, x_{p+1}) \\ &= \sum_{i=0}^{p+1} (-1)^i \varphi(x_0, x_1, \dots, x_{i-1}, \hat{x}_i, x_{i+1}, \dots, x_{p+1}) \end{aligned}$$

where “^” signs that we dropped the marked coordinate. Moreover, we proved the following properties

$$(1) \Phi_0^p(X, G) \subset \Phi^p(X, G)$$

$$(2) \Phi_0^p(X, G), \Phi_c^p(X, G)$$

$$(3) \delta(\Phi^p(X, G)) \subseteq \Phi_0^{p+1}(X, G)$$

$$(4) \delta(\Phi_c^p(X, G)) \subseteq \Phi_c^{p+1}(X, G)$$

for the sets

$$\Phi_0^p(X, G) = \{\varphi | \varphi \in \Phi^p(X, G), |\varphi| = \emptyset\} \text{ and}$$

$$\Phi_c^p(X, G) = \{\varphi | \varphi \in \Phi^p(X, G), |\varphi| = \text{compact}\}$$

and

$$\Phi_0^p(X, G) = \{\varphi | \varphi \in \Phi^p(X, G), |\varphi| = \emptyset\},$$

Where $|\varphi|$ is the *support of p-function*.

Lastly, we proved that the quotient spaces

$$C_c^p(X, G) = \Phi_c^p(X, G) / \Phi_0^p(X, G) \text{ and}$$

$$C^p(X, G) = \Phi^p(X, G) / \Phi_0^p(X, G)$$

inherit the properties of the Abelian mapping that a homeomorphism from the Abelian group $\Phi^p(X, G)$

2. Definition of Groups of Cohomologies

We define the sets

$$Z^p(X, G) = \{u | u \in C_c^p(X, G), \delta u = 0\}$$

$$B^p(X, G) = \delta[C_c^{p-1}(X, G)].$$

Since δ is a homomorphism between groups, it immediately follows that the sets $Z^p(X, G)$ and $B^p(X, G)$ are subgroups of the group $C^p(X, G)$. Moreover, these sets are actually the Kernel and the Image of the corresponding homomorphisms δ . They are called groups of **p-measurable cocycles** and **p-measurable coboundaries** (of topological space X with coefficients in the group G) correspondingly. It can be shown that for a homomorphism $\delta, \delta \circ \delta = 0$ is valid ([1]). By this we immediately have that the group of **p-measurable** coboundaries is a subgroup of the group of **p-measurable** cocycles $Z^p(X, G)$. This means that we can define the quotient group

$$H_c^p(X, G) = Z^p(X, G) / B^p(X, G)$$

This group is called **p -measurable group of cohomologies** with compact supports on topological space X , with coefficients in the group G . In this paper, we envisage the groups $H_c^p(X, G)$, for a locally compact topological space X . If we consider a compact topological space, then the group of cohomologies is denoted by $H_c^p(X, G)$. In such case, we also write $\Phi^p(X, G) = \Phi_c^p(X, G)$.

It is a general impression, at least to this level of presentation of definitions, that they are somehow unnatural and just scattered without any real meaning. However, as one digs deeper into the material, one discovers the importance and applications of theory of cohomologies, as well as it's not so apparent naturalness of the abovementioned definitions. Groups of cohomologies can be introduced, i.e., defined in lots of different ways. For compact topological spaces with „good” local behavior, different definitions lead to the same result. For „pathological” spaces however, different definitions can lead to the different results. Considering that each definition is based on a set of different assumptions they all have some advantages or/and drawbacks with respect to the other. In this presentation, we have chosen the approach that has the advantage of being general and yet simple. By this approach, we arrive very quickly to the subject of interest and study. In the following, we will present a very simple example of group of cohomologies.

3. An Example of Groups of Cohomologies

We notice that for any topological space X and any $p < 0$

$$H_c^p(X, G) = 0$$

Proposition 1: Let the topological space X consist of exactly one point, i.e., let $X = \{a\}$. Then

$$H_c^p(X, G) = 0, p > 0$$

and

$$H_c^0(X, G) \cong G$$

Lemma 1: Let the topological space X consist of exactly one point, i.e., let $X = \{a\}$. Then

1. $\Phi^p(X, G) \cong G$
2. $\Phi_0^p(X, G) = \{0\}$
3. $C_c^p(X, G) = C^p(X, G) = \Phi^p(X, G)$

Proof:

1. Let $p \geq 0$ be any given, but fixed natural number. Then the set of all $p+1$ -tuples (x_0, x_1, \dots, x_p) , where $x_i \in X$ and $i = 0, \dots, p$ has exactly one element (a, a, \dots, a) with a showing $p+1$ times. We can conclude that X^p is a one-element set. So, for any $p \geq 0$ and any p -function φ we can consider that

$$\varphi(x_0, x_1, \dots, x_p) = s \in G,$$

i.e., we can say that every element $\varphi \in \Phi^p(X, G)$ maps the $p+1$ -tuple (a, a, \dots, a) to a unique element $s \in G$. This way every element $\varphi \in \Phi^p(X, G)$ for any $p \geq 0$ determines a unique element of the group G . The opposite is also valid, i.e., each element of $s \in G$ determines a unique element $\varphi \in \Phi^p$. This means that we have a bijection between groups $\Phi^p(X, G)$ and G given by $f: G \rightarrow \Phi^p(X, G)$

$$f(s) = \varphi \Leftrightarrow \varphi(a, a, \dots, a) = s$$

We set $\varphi = \varphi_s$. The mapping f is also a homomorphism. Indeed, for any $s_1, s_2 \in G$ we have

$$\begin{aligned}
f(s_1 + s_2)(a, a, \dots, a) &= \varphi_{s_1 + s_2}(a, a, \dots, a) = s_1 + s_2 \\
&= \varphi_{s_1}(a, a, \dots, a) + \varphi_{s_2}(a, a, \dots, a) \\
&= (f(s_1) + f(s_2))(a, a, \dots, a).
\end{aligned}$$

This yields that the mapping f is an isomorphism, q.e.d.

2. Let $\varphi \in \Phi_0^p(X, G)$. Then the support $|\varphi| = \emptyset$ and for the element $a \in V$, $a \notin |\varphi|$. This means that there is a neighborhood V of $a, a \in V$, such that for every $(x_0, x_1, \dots, x_p) \in V$, $\varphi(x_0, x_1, \dots, x_p) = 0$. Actually, we have $\varphi(a, a, \dots, a) = 0$ with $X = \{a\}$. It follows that $\varphi = 0$. This shows that $\Phi_0^p(X, G) \subseteq \{0\}$. The opposite is always valid, so it follows that $\Phi_0^p(X, G) = \{0\}$.

3. It follows from 2, that

$$C^p(X, G) = \Phi^p(X, G) / \{0\} = \Phi^p(X, G) \cong G.$$

On the other hand, $\Phi_c^p(X, G) = \Phi^p(X, G)$, since a topological space of one point is always compact. It follows that

$$C_c^p(X, G) = \Phi_c^p(X, G) / \{0\} = \Phi^p(X, G) \cong G, \text{ q.e.d.}$$

We now proceed with the prove of the Proposition.

Let $p > 0$ be any natural number. Then, for any p -function $\varphi \in \Phi^p(X, G)$, we have

$$(\delta(\varphi))(a, a, \dots, a) = \sum_{i=0}^{p+1} (-1)^i \varphi(a, a, \dots, a, a, a, \dots, a),$$

where \hat{a} is on position with index i .

We will consider two cases:

1. Let $p > 0$ be an even number. Then, $p + 1$ is odd and on the right side we have even number of additions and so the sum will be zero. This means that

$$(\delta(\varphi))(a, a, \dots, a) = 0, \text{ i.e., } \delta(\varphi) = 0.$$

2. Let $p > 0$ be an odd number. Then, $p + 1$ is even and on the right side we have odd number of additions. This means that

$$(\delta(\varphi))(a, a, \dots, a) = -\varphi(a, a, \dots, a),$$

where on the left we have $p + 1$ -tuples and on the right we have p -tuples. The nature of the groups considered gives us

$$\delta(\varphi) = -\varphi,$$

because $\Phi_0^p(X, G) = \{0\}$ and $C_c^p(X, G) = \Phi^p(X, G)$.

As a result of the above we get:

1. For even $p > 0$

$\delta: C^p(X, G) \rightarrow C^{p+1}(X, G)$ is such that $\delta(\varphi) = 0$, for every $\varphi \in C^p(X, G)$ and we have that

$$Z^p(X, G) = C^p(X, G).$$

On the other side $p-1$ is odd and thus we have $\delta(\varphi) = -\varphi$, for every $\varphi \in C_c^{p+1}(X, G)$. It follows that

$$\delta[C_c^{p-1}(X, G)] = C_c^p(X, G),$$

i.e.,

$$B^p(X, G) = C_c^p(X, G).$$

Consequently we get

$$\begin{aligned} H_c^p(X, G) &= Z^p(X, G) / B^p(X, G) \\ &= C_c^p(X, G) / C_c^p(X, G) = \{0\}. \end{aligned}$$

2. For odd $p > 0$, we have that for every $\varphi \in C_c^{p+1}(X, G)$, $\delta(\varphi) = -\varphi$, i.e., we have

$$Z^p(X, G) = \{\varphi \in C_c^p(X, G), \mid \delta(\varphi) = -\varphi\} = \{0\}.$$

On the other side $p-1$ is even and thus we have $\delta(\varphi) = 0$, for every $\varphi \in C_c^{p-1}(X, G)$. It follows that

$$B^p(X, G) = \delta[C_c^{p-1}(X, G)] = \{0\}.$$

It follows that

$$H_c^p(X, G) = Z^p(X, G) / B^p(X, G) = \{0\} / \{0\} = \{0\}$$

If $p = 0$, then since it is considered an even number, for

$$\delta: C^p(X, G) \rightarrow C^1(X, G)$$

We have

$$Z^0(X, G) = C^p(X, G) = G$$

It follows that

$$H_c^0(X, G) = Z^0(X, G) / B^0(X, G) = G / \{0\} = G.$$

This proves the Proposition 1.

4. Conclusion

Following the definitions, structure and the general approach in [1], we presented a way to define groups of cohomologies. Then, in order to give life to these definitions, we've presented an example of such a group of cohomology. For our example we chosed a simple compact topological space of only one point. Following the sets and definitions we constructed the group of cohomologies $H_c^p(X, G)$ for this topological space. Moreover, we have given exact calculation of the coboundary operator δ . By doing this we are trying to give "life" to seemingly sterile and unnatural constructions.

References

- [1] Kotevska, E., Calamani, S. (2016). On some properties of groups of pcochains, *International Scientific Journal HORIZONS Series B* (2016), p. 7–15.
- [2] Kotevska, E. (1996). Cohomologies on a Locally Compact Topological Spaces, University St. *Cyril and Methodius*. Faculty of Natural Sciences, Institute of Mathematics .
- [3] McLane, S. (1963). Homology. Springer Verlag: Berlin.
- [4] Massey, W. (1978). Homology and Cohomology Theory. An Approach Based on Alexander-Spanier Cochains, M. Dekker (1978).
- [5] Neisendorfer, J. Primary homotopy theory, Mem. A.M.S., 232.
- [6] Notbohm, D. (1999). Spaces with polynomialmod- p cohomology. Mathematical Proceedings of the Cambridge Philosophical Society.

